



BURNSIDE

Swan Lake Aquatic Conditions Review

Friends of Swan Lake Park

**R.J. Burnside & Associates Limited
292 Speedvale Avenue West Unit 20
Guelph ON N1H 1C4 CANADA**

**October 2025 (rev Nov 2025)
300060899.0000**



Distribution List


No. of Hard Copies	PDF	Email	Organization Name
0	Yes	Yes	Friends of Swan Lake Park

Record of Revisions

Revision	Date	Description
-	October 31, 2025	Draft Submission to Friends of Swan Lake Park
1	November 7, 2025	Draft Submission to Friends of Swan Lake Park
2	November 26, 2025	Final Submission to Friends of Swan Lake Park

R.J. Burnside & Associates Limited

Report Prepared By:


 Mark Saunders, MES, CAN-CISEC-IT
 Aquatic Ecologist
 MS:af

Report Reviewed By:


 Chris Pfohl, C.E.T., EP, CAN-CISEC
 Senior Aquatic Ecologist

Executive Summary

- R.J. Burnside & Associates Limited (Burnside) reviewed aquatic habitat conditions in Swan Lake at the end of Phase 1 of the Long-Term Water Quality Management Plan, on behalf of the Friends of Swan Lake Park, to guide recommendations for Phase 2.
- High abundances of phytoplankton continue to cause turbidity and low dissolved oxygen, which may interfere with macrophyte re-establishment and the development of in-water habitat. Elevated internal nutrient loading and chloride concentrations may impede current and future restoration efforts if left unaddressed.
- Improving water circulation through a bioswale system and/or artificial aeration may help reduce internal nutrient loading could assist in nutrient management by helping to remove and sequester internal phosphorus loading.
- High chloride levels, limited structural habitat, and predation from abundant Fathead Minnow populations may be suppressing zooplankton grazers (e.g., *Daphnia* spp.), although no study has yet confirmed this relationship.
- Largemouth Bass stocking in 2025 provides an opportunity to potentially reduce Fathead Minnows. Establishment of a population of bass may be supported by adding complex woody material structures in the absence of macrophytes. Future aquatic surveys are recommended to determine whether current lake conditions are suitable for sustaining bass, especially to over winter, or if further habitat modifications (e.g., deepening the pond or raising the lake level) may be worth exploring to improve survivability. However, it is worth evaluating the bass survival in 2026 to see if these deepening measures are necessary.
- There are limited opportunities to use native Unionidae mussels or sunfish species to improve habitat quality.
- Due to high turbidity from phytoplankton preventing macrophyte growth, efforts should focus on improving water quality before pursuing further macrophyte restoration.

Table of Contents

Distribution List.....i
Record of Revisions.....i
Executive Summaryii
Table of Contentsiii
1.0 Background.....1
2.0 Literature Review Methods.....2
3.0 General Overview of End of Phase 13
4.0 Turbidity and Dissolved Oxygen.....6
 4.1 Artificial Aeration.....6
 4.2 Evaluation of Proposed Bioswale.....8
5.0 Chloride10
6.0 Zooplankton and Macroinvertebrates11
 6.1 Zooplankton (*Daphnia* spp.).....11
 6.2 Macroinvertebrates11
 6.3 Freshwater Mussels.....12
7.0 Fish Community.....13
 7.1 Control of Minnows via Biomanipulation13
 7.2 Limitations to Largemouth Bass Survival in Swan Lake14
 7.3 Other Fish as a Means of Biomanipulation.....16
8.0 Aquatic Vegetation17
9.0 Recommendations.....19
10.0 Conclusion21
1.0 References22

Tables

Table 1: Interim Water Quality Targets for the Phase 1 (2021-2025) Adapted from FOSLP (2025a)3
 Table 2: Average Inorganic Water Parameters (Mean ± SD) in Swan Lake during 2025 (April 24-Sept 24) n=1-6.4
 Table 3: Summary of Aeration Recommendations for Swan Lake from Anderson and Baldwin (2025).7
 Table 4: Lake Turnover (80,000 m³) from April-October (214 days). Adapted from FOSLP (2023).9
 Table 5: TRCA Swan Lake Fish Surveys During Phase 1. Bottom Feeding Fish (Catfish, Common Carp, and Goldfish) Were Euthanized. All Other Species Returned Alive to the Lake (FOSLP 2025b)..... 14
 Table 6: Macrophyte Types Observed in Swan Lake 1993 and 2024. Adapted from FOSLP (2025b) 18

Table 7: Summary of Recommendations to Improve Biological/Ecological Conditions
During Phase 2.....19

Figures

Figure 1: Swan Lake and Runoff Monitoring Stations. Image provided by FOSLP. Image Quality Improved through GPT-5 (Open AI 2025a).....1

Figure 2: Proposed Route for Bioswales. Image from FOSLP (2023). Image Quality Improved through GPT-5.....8

Figure 3: Location of Three Oil Grit Separators (Red Circles) Responsible for 56% of Chloride Inputs. From FOSLP (2023).10

Figure 4: Summary of Swan Lake Macroinvertebrate Survey (2023). Image Modified from FOSLP (2025b)12

Figure 5: Proposed Biomanipulation to Achieve a Top-Down Trophic Cascade in Swan Lake. Created with assistance from GPT-5 (OpenAI 2025b).....13

Figure 6: Bathymetric Map of Swan Lake. Provided by FOSLP.15

Photos

Photo 1: North Channel Dry at Time of Inspection, Containing Remnants of Landfill Waste. Represents a Beautification and Bioswale Opportunity.....9

Swan Lake Aquatic Conditions Review
October 2025 (rev Nov 2025)

List of Abbreviations

DO – Dissolved Oxygen

FOSLP – Friends of Swan Lake Park

TRCA – Toronto and Region Conservation Authority

SWM – Stormwater Management

OGS – Oil Grit Separator

PAC – Poly-aluminum Chloride

Disclaimer

Other than by the addressee, copying or distribution of this document, in whole or in part, is not permitted without the express written consent of R.J. Burnside & Associates Limited.


1.0 Background

Swan Lake is a 5.5 ha artificial lake in the City of Markham, created from a former gravel quarry excavated during the 1960s–1970s (FOSLP 2025a). Once dewatering operations ceased, the pit filled naturally with groundwater. During the 1970s, the site was used as a local landfill, and some illegal dumping occurred.

By the 1990s, the area surrounding Swan Lake was slated for urban development (FOSLP 2025a). The *Swan Lake Community Environmental Management Study* (December 1993) proposed converting the quarry into a public park and community feature with restored aquatic and terrestrial habitat. Markham accepted the plan in 2000.

Swan Lake has no natural surface inflows or outflows (FOSLP 2025a; Figure 1). It was incorporated into the stormwater management (SWM) system, receiving runoff from six stormwater sources and discharging through one outflow that regulates water levels at ~208.3 m ASL, with an average depth of 1.65 m.

Figure 1: Swan Lake and Runoff Monitoring Stations. Image provided by FOSLP. Image Quality Improved through GPT-5 (Open AI 2025a)



Swan Lake Aquatic Conditions Review
October 2025 (rev Nov 2025)

Persistent poor water quality and recurring algal blooms led local residents to form the Friends of Swan Lake Park (FOSLP) in 2019. Their goal is the rehabilitation of aquatic and terrestrial habitats to achieve sustainable water quality for long-term human and wildlife use. In 2020 FOSLP submitted *A Pathway to Sustainable Water Quality for Swan Lake* to City Council, prompting the 25-year *Long-Term Water Quality Management Plan* launched in 2021. The plan's stated goal is "to improve the overall health of Swan Lake, which will provide opportunities for no-contact activities for the enjoyment of the community" (FOSLP 2025a).

Thus far, Phase 1 has been completed and now progressing into Phase 2:

- **Phase 1 (2021-2025)** targeted immediate water-quality improvements (chemical phosphorus reduction, geese management, invasive-species management, and pilot aquatic plantings)
- **Phase 2 (2026-2030)** is under review by Markham. FOSLP is recommending improved oxygen levels through circulation improvements, chloride reduction, sustaining nutrient levels and stronger ecological monitoring

FOSLP retained R.J. Burnside & Associates Limited (Burnside) to provide a preliminary review of aquatic habitat conditions at the end of Phase 1 and provide recommendations entering Phase 2. The following focuses on impacts and recommendations to support potential improvements for aquatic life (zooplankton, invertebrates, fish, and macrophytes).

2.0 Literature Review Methods

A literature review was conducted of the most recent reports commissioned in relation to the Swan Lake study, including those previously prepared for or by the FOSLP. Burnside visited Swan Lake with Fred Peters of FOSLP on October 15, 2025, to observe the condition of the lake, stormwater management facilities, and surrounding parklands.

Additional literature was reviewed on small, shallow, eutrophic, temperate lakes, with an emphasis on studies from North America or the northern hemisphere. While recent publications (post-2010) were prioritized, foundational and frequently cited papers were also referenced to provide context and theoretical background.

Portions of this report were edited using AI-assisted editing tools in ChatGPT (OpenAI) to improve clarity and readability. AI was also used to help improve poor figure quality and to generate an illustrative figure (OpenAI, 2025a, 2025b). All research, technical analysis, writing, and interpretations are the author's own.

3.0 General Overview of End of Phase 1

Throughout Phase 1, total phosphorus and nitrogen species levels have improved from hypereutrophic conditions in 2020 to mesotrophic conditions by 2024 (FOSLP 2025a). The most recent FOSLP report indicates that the interim water-quality targets for Phase 1 were nearly achieved (Table 1). Monitoring data collected in 2025 support the continued positive trends observed during Phase 1 (Table 2). Nutrient concentrations (i.e., total phosphorus and nitrogen species) remain low and below target thresholds, indicating ongoing improvement in water quality. However, chloride concentrations and conductivity remain elevated (approximately 200 mg/L).

Table 1: Interim Water Quality Targets for the Phase 1 (2021-2025) Adapted from FOSLP (2025a)

Parameter	Current Value	Interim Target	Objective and Rationale
Total Phosphorus (mg/L)	>0.2	0.05-0.1	Current Value: The average of the growing season TP values in the period since 2016 has been 0.2 mg/L. Interim target: will provide a low eutrophic condition in the first year after treatment increasing to eutrophic in year three.
Secchi Transparency (m)	<0.5	0.6-0.8	Based on correlation with the phosphorus target. Secchi is also a substitute for Chlorophyll a.
Frequency of Algae Blooms	Annual	Every Three Years	Trigger for treatment every three years.
Internal Phosphorus Load (kg/yr)	53	0-25	Both internal and external loads should be controlled to achieve the lake conservation target (see above)
External Phosphorus Load (kg/yr)	30	15	

Table 2: Average Inorganic Water Parameters (Mean ± SD) in Swan Lake during 2025 (April 24-Sept 24) n=1-6.

Parameter	Surface (S2025)	Dock 0.5m (S105)	Dock 1.5 m (S115)	Mid Lake (Ultrasound Transducer)	Trophic State
Total P (mg/L)	0.018±0.008	0.013±0.004	0.016±0.005	0.021±0.002	Eutrophic (0.031-0.100) – below interim target
Orthophosphate (mg/L)	0.006	ND	ND	ND	Minimal soluble P
Nitrate + Nitrite (m/L)	ND	ND	ND	0.11	Minimal Nitrogen species.
Total Kjeldahl Nitrogen (mg /L)	0.59±0.13	0.56±0.05	0.56±0.10	0.53±0.06	Stable, moderate nutrient pool
Ammonia (µg /L)	0.130	0.072±0.023	0.120±0.028	0.070±0.009	Small fraction of total ammonia is the toxic un-ionized NH ₃ at average Lake pH ≈ 8.1–8.2 (CCME 2010).
Chloride (mg/L)	213±17.1	217±17.5	200	190	Above >120 mg/L guidelines (CCME 2011)
Conductivity (µS/cm)	907±65	940±69	930	920	Elevated from salt.
Dissolved Organic Carbon (mg/L)	8.5±0.96	8.3±0.88	-	5.4±0.50	Moderate organic matter
pH	8.1±0.27	8.2±0.28	-	8.1±0.01	Normal, slightly basic
ND = Not Detected at a concentration equal or greater than the indicated Detection Limit of 0.004 mg/L for Orthophosphate and 0.1 mg/L for Nitrate + Nitrite. If a value appears without a ± SD, it indicates that dataset consisted of only one observation (n=1) or repeated identical observations					

The improvements in water quality can be attributed to nutrient control measures enacted in Phase 1. Chemical treatment has been the main phosphorous control throughout the lake's history. Before the *Management Plan*, Copper-sulfate treatment was applied 1995, and a 25.2 t Phoslock® application in 2013 (FOSLP 2025a). Under Phase 1, Markham switched to poly-aluminum chloride (PAC), with a 13 t application in 2021 and 9.4 t application in 2024 (~\$150,000 per application). While effective in the short-term, this approach treats symptoms rather than sources.

External nutrient reduction has focused on Canada goose hazing and shoreline fencing to limit fecal loading (FOSLP 2025a). The underlying aquifer appears to facilitate limited flushing of nutrient-rich water, gradually improving chemistry.

Despite nutrient reductions, turbidity caused by phytoplankton (i.e., algae in the water column) overgrowth and bottom-water anoxia (0 mg/L dissolved oxygen (DO)) continues to persist. To address phytoplankton-related turbidity, a Water IQ Pulsar 4400+ ultrasonic unit was installed in 2024; however, turbidity remains largely unchanged (FOSLP 2025a). Frequent phytoplankton blooms also contribute to anoxia as decomposing organic material consumes dissolved oxygen. Continuous DO monitoring began at the dock in 2023 (FOSLP 2025a). By 2024, surface oxygen levels remained adequate (>5 mg/L), but anoxic conditions persisted near the lake bottom during the summer.

Markham and FOSLP have diverging opinions on how to interpret the water quality results of Phase 1:

City of Markham / AECOM Position

- View surface-oxygen improvement and nutrient reduction as successes; believe nutrient control alone will maintain DO.

FOSLP Position

- Argues improvements are temporary; Swan Lake's artificial structure requires active oxygenation or circulation.
- Proposed calcium-peroxide dosing and mechanical oxygenation pilots (i.e., Fleming College 2022; University of Toronto 2023) were rejected for "unproven" or "sediment-disturbance" concerns (FOSLP 2025a).
- Maintains that controlled oxygenation could stabilize redox conditions and reduce internal P release.

In the following sections, we discuss the ecological consideration for aquatic life that could inform possible actions taken over the course of Phase 2.

4.0 Turbidity and Dissolved Oxygen

4.1 Artificial Aeration

FOSLP commissioned Natural Resource Solutions Inc. (Anderson and Baldwin 2025) to review current oxygenation management strategies for Swan Lake. The resulting recommendations are summarized in Table 3. Overall, the report concludes that as Swan Lake recovers, its biological oxygen demand (BOD) will likely increase as aerobic organisms (e.g., bacteria, fish, and benthic invertebrates) become more active and abundant. Consequently, any aeration system must be capable of supporting both current and future BOD as the ecosystem responds to rehabilitation efforts. The report determined that diffused aeration and fountain aeration are the most suitable methods for improving oxygenation and circulation within the lake.

As detailed in the report (see reference within Anderson and Baldwin 2025), improving DO levels may not only address the anoxia observed in the deeper portions of the lake (creating conditions that allow more characteristic aquatic life to thrive) but also help reduce the high levels of turbidity currently observed. In the presence of oxygen, phosphorus in the water binds to iron and clay particles and settles to the lake bottom, reducing its availability to phytoplankton in the water column and thereby limiting phytoplankton growth. Overtime, less phytoplankton in the water column should result in clearer water.

Table 3: Summary of Aeration Recommendations for Swan Lake from Anderson and Baldwin (2025).

Method	Mechanism of Action	Advantages	Limitations	Feasibility for Swan Lake
Diffused Aeration	Agitates or circulates water at the surface to enhance gas exchange and disrupt localized stagnation. Releases compressed air through submerged diffusers to promote vertical mixing and oxygen transfer.	Immediate DO improvement; simple installation; visible circulation. Gentle, uniform oxygenation; low disturbance; energy-efficient; adaptable to seasonal use.	Limited spatial reach; energy-intensive; risk of sediment resuspension. Requires careful diffuser placement; effectiveness reduced in very shallow areas.	Highly suitable; offers broad coverage and sustained oxygenation when operated continuously.
Fountain Aeration	Pumping water through surface-level diffusers into the air to promote gas exchange and surface agitation.	Uniform oxygenation; low disturbance; simple installation.	Limited spatial reach; requires careful fountain placement; non-functional during winter months.	Highly suitable; simple installation; offers sustained oxygenation when operated continuously.
Chemical Oxygenation	Introduces oxygen-releasing or oxidizing compounds that supply elemental oxygen or modify sediment redox conditions.	Direct oxygen delivery; long-lasting effect; no mechanical disturbance.	Requires precise dosing; limited circulation benefit; may provide only temporary improvement.	Moderate feasibility; useful as supplemental treatment for localized low-oxygen zones.
Destratification / Mixing	Induces circulation to break down stratification and promote natural oxygenation through gas exchange.	Reduces stagnation; minimizes hypoxic zone formation; enhances natural mixing.	Limited DO increase; potential warming; continuous operation required.	Partially feasible; could aid circulation but unlikely to raise overall DO significantly.

Despite the promise of mechanical aeration, recent studies suggest that such actions are not a “silver bullet.” Artificial aeration in shallow, temperate lakes often has only modest effects in preventing further phosphorus release (May et al. 2020) and can be ineffective in systems with large surface areas (Kibuye et al. 2021), such as Swan Lake. Therefore, mechanical aeration may be most effective if used in the deeper reaches of the lake. Incorporating additional measures to reduce internal nutrient loading would likely enhance the effectiveness of any oxygenation program, though it alone is unlikely to result in water quality improvements.

4.2 Evaluation of Proposed Bioswale

High internal phosphorus loading is a persistent challenge in shallow lake recovery (Søndergaard et al. 2003; May et al. 2020; Kibuye et al. 2021). While creating oxygenated conditions typically suppresses phosphorus availability, shallow lakes function differently than deeper systems because the distance between the sediment surface (where phosphorus is stored) and the photic zone (where phytoplankton grow) is short (Søndergaard et al. 2003). As a result, it takes little energy for nutrients to become resuspended. Wind events or the decomposition of settled organic material can quickly reintroduce phosphorus into the water column, where it becomes available for phytoplankton growth (Søndergaard et al. 2003).

In the case of Swan Lake, internal loading appears to be a more significant concern than external loading (FOSLP 2025a). With assistance from Fleming College (Siembida-Lösch 2021), FOSLP (2023) proposed investigating bioswale options, particularly within the 100 m North Channel and the Turtle Inlet (Figure 2), as a means of reducing internal nutrient loads. Siembida-Lösch (2021) provides an extensive review of bioswale effectiveness, and a more

recent study emphasizes that design should prioritize nutrient removal efficiency (Ali and Pickering 2022).

Figure 2: Proposed Route for Bioswales. Image from FOSLP (2023). Image Quality Improved through GPT-5 (Open AI 2025a)



Design outlines provided by Siembida-Lösch (2021) and FOSLP (2023) emphasize the importance of achieving a hydraulic detention time (HDT) of < 30 days to control phytoplankton growth. To meet this target for Swan Lake (80,000-102,000 m³), Siembida-Lösch (2021) recommended recirculating approximately 3,400 m³ per day. However, lower rates proposed by FOSLP (2023) could provide lower-cost alternatives while still

achieving desirable outcomes, with up to 27% of the lake volume recycled (Table 4). Smaller pumps could be powered by existing on-site renewable electricity or by restoring the wind turbines currently present on-site.

Table 4: Lake Turnover (80,000 m³) from April-October (214 days). Adapted from FOSLP (2023).

Pump Size	Capacity (m ³ /24h)	% of Lake Volume per Day	% of Lake Volume (50% Operation)
1 HP	50	13%	7%
2 HP	200	54%	27%

It should be noted that properly designed bioswales can effectively remove particulate phosphorus but may struggle to remove dissolved phosphorus from the water column (Ali and Pickering 2022). Fortunately, dissolved phosphorus concentrations in Swan Lake appear to be very low as of 2025 (Table 2). The addition of biochar, recommended by York University as a means of aiding nutrient and chloride filtration (FOSLP 2025a); however, recent studies have demonstrated variable effectiveness at removing nutrients (Ali and Pickering 2022).

The underused North Channel, which is currently dry, may represent the most suitable location for an initial bioswale pilot, as it would minimize disturbance to the Turtle Inlet habitat. In addition, because the North Channel contains remnants of the former landfill (Photo 1),

Photo 1: North Channel Dry at Time of Inspection, Containing Remnants of Landfill Waste. Represents a Beautification and Bioswale Opportunity



selecting this site could also present an opportunity to beautify and ecologically restore this area within the existing channel.

As noted by Kibuye et al. (2021), reducing external nutrient loading is the first and most critical step in achieving sustained reductions in phytoplankton blooms. Even if a bioswale is constructed, ensuring a reduction (or current levels) of nutrient inputs will help. Migratory geese are identified as the largest contributors to external nutrient inputs into the lake (FOSLP 2025a). Geese hazing has proven effective and may be the only feasible action to deter migratory geese in an


urban environment, as shoreline plantings (i.e., replacing manicured lawns with forests) are mainly targeted for residential geese.

5.0 Chloride

Chloride levels have significantly improved over Phase 1, declining from over 700 mg/L in 2020 to approximately 200 mg/L by 2024; a reduction of more than 70 % (FOSLP 2025a). No active measures were taken to achieve this reduction in chlorides, which is likely a result from clearing blockages at the east SWM Pond (#105) that had increased flow bypassing the pond and exchange with the underlying aquifer, combined with minimal surface outflow during the summers of 2022-2023 (FOSLP 2025a).

Three Oil Grit Separator (OGS) units have been identified as point-source contributors responsible for roughly 56% of chloride inputs (Figure 3), as they discharge directly into the lake rather than first passing through SWM ponds (FOSLP 2025a). Debate remains whether to

Figure 3: Location of Three Oil Grit Separators (Red Circles) Responsible for 56% of Chloride Inputs. From FOSLP (2023).



eliminate these discharges entirely by rerouting the stormwater at a cost estimated at \$1.8 million or retrofit filtration systems to intercept inflows. Engineering and financial factors will guide that decision.

At the current chloride concentration (~200 mg/L), further ecological improvement is unlikely. Persistent chloride enrichment may be sustaining high phytoplankton levels by suppressing zooplankton grazers that normally control phytoplankton (Hintz et al. 2022). The following section examines this theory and related aquatic ecosystem issues.

6.0 Zooplankton and Macroinvertebrates

Often overlooked in favor of more charismatic species such as sport fishes (e.g., bass and sunfish), zooplankton, macroinvertebrates (e.g., aquatic insects, worms, and crustaceans visible with naked eye), and freshwater mussels form the base of the freshwater food web (Hintz et al. 2022), supporting fish, amphibians, and birds. Ontario has few native obligate herbivorous or planktivorous fish; thus, zooplankton, macroinvertebrates, and mussels act as the principal phytoplankton / algae grazers in these systems (Newman 1991; Sarnelle 1992). Consequently, maintaining healthy communities of these taxa may be critical for phytoplankton control.

6.1 Zooplankton (*Daphnia* spp.)

Zooplankton are more sensitive to chloride than fish (CCME 2011; Hintz et al. 2022). The current chloride concentration (~200 mg/L) is below levels harmful to most fishes but falls within, or slightly above, the chronic exposure range for key grazers such as water fleas in the *Daphnia* genus, reducing phytoplankton/algal grazing capacity (CCME 2011; Hintz et al. 2022). Even at the 120 mg/L federal target, some *Daphnia* spp. populations show reduced survival and reproduction (Hintz et al. 2022; Woodley et al. 2023).

Zooplankton, especially *Daphnia* spp., are the dominant grazers of phytoplankton in temperate freshwater lakes (Sarnelle 1992). They can help regulate cyanobacteria through grazing but do not provide a reliable or sustained means of cyanobacteria bloom elimination (May et al. 2020; Kibuye et al. 2021). Thus, *Daphnia* spp. may be uniquely suited to aid in the recovery of Swan Lake directly in a way no fish species likely could. FOSLP (2025b) has already noted the possibility of chloride-driven suppression of zooplankton, though no formal studies of Swan Lake have been completed to confirm this pattern.


Daphnia spp. stocking has been demonstrated as an effective strategy in other eutrophic temperate lakes when implemented in tandem with the introduction of piscivorous fish (see Section 7; Ha et al. 2012). However, other studies have shown that natural recovery of zooplankton communities can occur when environmental conditions improve (Albright et al. 2004; May et al. 2020; Kibuye et al. 2021). Since there is no current data available on the existing zooplankton community composition (i.e., whether it is dominated by inefficient grazers such as rotifers or includes effective grazers such as *Daphnia* spp.), *Daphnia* spp. stocking is not recommended at this time. Instead, assessing the present zooplankton community is advised to determine whether the working hypothesis (i.e., low *Daphnia* spp. abundance may be contributing to elevated phytoplankton biomass) holds true.

6.2 Macroinvertebrates

Macroinvertebrates, while important benthic phytoplankton/algal grazers (Newman 1991), are strongly influenced by eutrophication but do not directly mitigate it (May et al. 2020). Many macroinvertebrates are more resilient than zooplankton and are unlikely to be severely affected by current chloride levels (CCME 2011). Instead, in a system such as Swan Lake, the

macroinvertebrate composition reflects improvements in oxygen, sediment quality, and habitat during recovery (May et al. 2020). Although FOSLP requested a macroinvertebrate study, the City of Markham declined (FOSLP 2025a). In 2023, FOSLP independently commissioned Mr. Chris Reeves to conduct a macroinvertebrate survey (FOSLP 2025b). The results indicated that Swan Lake supports a fair-to-intermediate water-quality community, with reduced diversity near stormwater inflows (Figure 4), suggesting that poor water quality continues to shape community structure (FOSLP 2025b).

Figure 4: Summary of Swan Lake Macroinvertebrate Survey (2023). Image Modified from FOSLP (2025b)



6.3 Freshwater Mussels


The use of freshwater mussels to control phytoplankton may be limited in Swan Lake. Although mussels have been shown to reduce phytoplankton / algal biomass, most demonstrations have involved the invasive Zebra Mussel (*Dreissena polymorpha*), making such applications undesirable (Geletu 2022). Native Unionidae mussels in Ontario have limited tolerance to polluted waters and possess complex reproductive strategies that require specific host fish (Metcalf-Smith et al. 1998). As a result, they are generally poor colonizers of new habitats. Many species are also restricted by chloride sensitivity during their larval (glochidia) stage, with reductions to survival (EC₁₀) ranging from 24 to 789 mg/L, while adults may tolerate chloride concentrations near 200 mg/L (CCME 2011). While native floater mussels (subfamily Anodontinae) tend to be more pollution-tolerant, the Paper Pondshell (*Utterbackia imbecillis*) may represent a potential candidate species, as it is more tolerant of degraded water quality, inhabits shallow ponds, and uses readily available host fish such as Largemouth Bass (*Micropterus salmoides*) and sunfishes (*Lepomis* spp.) (Keller and Ruessler 1997; Metcalf-Smith et al. 1998).

7.0 Fish Community

7.1 Control of Minnows via Biomanipulation

In Swan Lake, high populations of species such as Fathead Minnow (*Pimephales promelas*) may contribute to reduced zooplankton abundance and weakened phytoplankton control within the water column, consistent with a trophic cascade dynamic (Power 1992). In addition, in shallow lakes such as Swan Lake, zooplankton are more exposed to predation and thus exert weaker grazing pressure on phytoplankton than deeper systems that provide refuge zones (May et al. 2020). Adequate macrophyte coverage can offer such refuge and restore balance (Kibuye et al. 2021), but the current turbid conditions and high chlorides limit macrophyte recovery (see Section 8). Thus, Swan Lake currently offers minimal physical habitat conducive to the recovery of zooplankton even in the absence of high chlorides.

Figure 5: Proposed Biomanipulation to Achieve a Top-Down Trophic Cascade in Swan Lake. Created with assistance from GPT-5 (OpenAI 2025b).



Introducing or enhancing piscivore populations such as Largemouth Bass, could help to improve water quality through biomanipulation via top-down trophic control (Carpenter et al. 1985; May et al. 2020; see Figure 5). A classic example of such biomanipulation is described by Albright et al. (2004), who studied a small, shallow eutrophic pond (15.6 ha, mean depth 1.8 m) in upstate New York that, prior to intervention, contained only minnows (e.g., Golden Shiner, *Notemigonus crysoleucas*) and Brown Bullhead (*Ameiurus nebulosus*). The lake exhibited high phytoplankton biomass, elevated turbidity, and a zooplankton community of low diversity, dominated by rotifers and few *Daphnia* spp. In 1999, Largemouth

Introducing or enhancing piscivore populations such as Largemouth Bass, could help to improve water quality through biomanipulation via top-down trophic control (Carpenter et al. 1985; May et al. 2020; see Figure 5). A classic example of such biomanipulation is described by Albright et al. (2004), who studied a small, shallow eutrophic pond (15.6 ha, mean depth 1.8 m) in upstate New York that, prior to intervention, contained only minnows (e.g., Golden Shiner, *Notemigonus crysoleucas*) and Brown Bullhead (*Ameiurus nebulosus*). The lake exhibited high phytoplankton biomass, elevated turbidity, and a zooplankton community of low diversity, dominated by rotifers and few *Daphnia* spp. In 1999, Largemouth

Bass and Smallmouth Bass (*Micropterus dolomieu*) were introduced, and by 2002 water quality visibly improved as minnow populations declined and *Daphnia* spp. increased in abundance and diversity. However, controlling planktivorous fish populations does not always yield such clear results; some studies report that biomanipulation has little or no measurable effect on zooplankton communities (Setubal and Riccardi 2020).

Largemouth Bass are currently stocked into the lake as of 2025 (FOSLP 2025a). Bass were also stocked 1992, but fish surveys conducted by the Toronto Region Conservation Authority (TRCA) do not seem to list Largemouth Bass among species capture (Table 5).

Table 5: TRCA Swan Lake Fish Surveys During Phase 1. Bottom Feeding Fish (Catfish, Common Carp, and Goldfish) Were Euthanized. All Other Species Returned Alive to the Lake (FOSLP 2025b)


Date	Fish Species	Number of Fish
April 2021 (three days electrofishing + two days nets)	Brown Bullhead	210
	Common Carp	7
	Fathead Minnow	>10,000
April 2022 (one day electrofishing + one day nets)	Brown Bullhead	80
	Common Carp	20
	Fathead Minnow	875
April 2023 (one day electrofishing)	Brown Bullhead	84
	Common Carp	103
	Fathead Minnow	14
	Goldfish	2
April 2024 (one day electrofishing + one day nets)	Brown Bullhead	193
	Common Carp	1
	Fathead Minnow	1,521
	Goldfish	13
	Common Carp X Goldfish	59
	Emeral Shiner	1

7.2 Limitations to Largemouth Bass Survival in Swan Lake

The shallow morphometry of Swan Lake, with its broad surface area and limited depth, may constrain the successful establishment of Largemouth Bass. Although Largemouth Bass occupy a wide range of depths, from very shallow (<1 m) to deeper water (~12 m) (Peat et al. 2016), they typically prefer depths of 2-3 m., especially in winter (Bunt et al. 2021). Such depths are scarce in Swan Lake, where nearly 60% of the basin is less than 1 m deep (Figure 6). Shallow lakes are especially vulnerable to meteorological variability and experience rapid temperature fluctuations in response to changes in air temperature (Piccioni et al. 2021), creating thermal instability that challenges overwintering bass. Feeding activity ceases below 6°C, making

Largemouth Bass sensitive to prolonged cold periods (Fullerton et al. 2000; Bunt et al. 2021). DO is another critical factor for overwinter survival, ideally remaining within the 2-6 mg/L range (Hasler et al. 2009). Current DO measurements are limited to nearshore summer readings, but values at ~2 m depth suggest potential for winter anoxia.

Figure 6: Bathymetric Map of Swan Lake. Provided by FOSLP.



Bass stocking efforts should prioritize introducing larger individuals (>100 mm total length) to improve survival and establishment (Fullerton et al. 2000; Post et al. 1998). Bass strongly prefer in-water structure provided by macrophyte beds and coarse woody material (Bunt et al. 2021). Field observations indicate that Swan Lake’s littoral zone is dominated by short meadows of invasive Eurasian Milfoil (*Myriophyllum spicatum*), with few other macrophytes and a largely featureless limnetic zone. Installing coarse woody material (e.g., upright stumps with root mass, properly elevated logs and branches) could provide immediate cover and overwintering refuge habitat, that can be locally sourced until native vegetation recovers (Czarnecka 2016). TRCA has proposed shoreline restoration efforts that include addition of woody material and rock complexes in the littoral zone (TRCA 2022). Thus, such efforts have already been under consideration in the case of Swan Lake. Consideration of placing deeper water structures (stumps and log cover) is also required to provide refuge from predatory birds and human anglers.

Deepening portions of Swan Lake could create refuge pools suitable for overwintering Largemouth Bass, although suitable water quality needs to be confirmed during all seasons. Such modifications may also align with community goals to enhance recreational opportunities around the lake. Alternatively, modestly increasing the lake's regulated water level could expand available overwintering habitat and would be less expensive than dredging. Currently, approximately 12% of Swan Lake offers adequate depth for overwintering, assuming the lake does not become anoxic.

Future aquatic surveys will confirm whether Largemouth Bass can successfully overwinter under current lake conditions and help determine if habitat enhancements are warranted. If Largemouth Bass are detected in the 2026 surveys, continued stocking of larger individuals (>100 mm) is recommended, along with the placement of woody material complexes (stumps with root masses and elevated log structures) to provide habitat until macrophyte recovery occurs. Re-evaluation of the need for more extensive interventions (e.g., dredging) should be deferred until bass survival is confirmed, as the current stocking provides an opportunity to assess the effectiveness of existing conditions before pursuing additional modifications.

7.3 Other Fish as a Means of Biomanipulation

As previously noted, there are few planktivorous fish that could be introduced to control phytoplankton growth suitable to Ontario, with the most common species being the highly invasive Bighead Carp (*Hypophthalmichthys nobilis*) and Silver Carp (*Hypophthalmichthys molitrix*), which are undesirable species in most circumstances (Geletu 2022). One of the few native planktivorous fish in Ontario is the Gizzard Shad (*Dorosoma cepedianum*), but its effectiveness in controlling phytoplankton remains unclear. Although Gizzard Shad feed on phytoplankton, their high waste excretion rates rapidly recycle nutrients back into the water column, often stimulating rather than reducing algal growth (Schaus et al. 2010). In fact, removal of Gizzard Shad has been demonstrated as an effective means to reduce phytoplankton levels (Godwin et al. 2011).

Bluegill (*Lepomis macrochirus*) were also considered for stocking in Swan Lake by TRCA (FOSLP 2025a); however, smaller sunfishes such as Bluegill may not be desirable for improving the water quality of Swan Lake. Sunfishes are important zooplankton feeders in North American lakes and may therefore contribute to increased phytoplankton biomass by preferentially consuming *Daphnia* spp. or other zooplankton such as rotifers (Noda and Maruyama 2013; Bernes et al. 2015). Accordingly, the addition of sunfish species is not recommended at this stage, as it may exacerbate the elevated phytoplankton levels currently observed in Swan Lake.

8.0 Aquatic Vegetation

Macrophytes are effective at stabilizing sediments, improving water clarity, and reducing nutrient concentrations (Kibuye et al. 2021). However, the current turbid conditions in Swan Lake likely prevent their establishment, limiting these benefits. Previous macrophyte plantings have failed under the lake's turbid, nutrient-rich, and high-chloride conditions. Plantings of Wild Celery (*Vallisneria americana*) in 2023 and 2024 experienced >90% mortality (FOSLP 2025b), primarily due to light limitation and turbidity. Considering the limited success of past plantings and supporting literature, further macrophyte restoration efforts are not recommended until measurable improvements in water clarity occur.

Submerged macrophytes, not rigid emergent species such as reeds or cattails, provide superior habitat complexity for Largemouth Bass and other fish, supporting higher growth and recruitment at moderate coverage (~60 %) (Stahr and Shoup 2015; Nohner et al. 2018). Future planting should therefore focus on submerged and floating-leaved species, rather than emergent vegetation, to maximize habitat value and promote long-term ecological balance (Table 6). In fact, macrophyte recovery in similarly sized eutrophic lakes has been linked to the introduction of piscivorous fish (Albright et al. 2004). Thus, these efforts are likely to synergize.

While macrophyte beds can reduce nutrient concentrations (particularly phosphorus) by absorbing nutrients, stabilizing sediments, and increasing oxygen near the substrate, they can also, under certain conditions, contribute to internal phosphorus loading. Dense vegetation may create localized anoxia and elevated pH, both of which can enhance phosphorus release from sediments (Søndergaard et al. 2003). This potential effect should be monitored as vegetation is re-established.

Table 6: Macrophyte Types Observed in Swan Lake 1993 and 2024. Adapted from FOSLP (2025b)

Open Waters: 0.5 m to 1.5 m Deep		Identified in 1993	Identified in 2024	Recommended
Pond Lily	<i>Nuphar variegata</i>			Yes
Water Lily	<i>Nymphaea odorata</i>			Yes
Pondweed	<i>Potamogeton richardsonii</i>			Yes
Pondweed	<i>Potamogeton pectinatus</i>			Yes
Lake Margins and Water to 0.75 m Deep		Identified in 1993	Identified in 2024	Recommended
Bur-reed	<i>Sparganium eurycarpum</i>			
Sweet Flag	<i>Acorus calamus</i>			
Soft-stem Bulrush	<i>Schoenoplectus tabernaemontani</i> (formerly <i>Scirpus validus</i>)			
Shores and Waters to 0.5 m Deep		Identified in 1993	Identified in 2024	Recommended
Narrow-leaved Cattail	<i>Typha angustifolia</i>	X	X	
Wide-leaved Cattail	<i>Typha latifolia</i>		X	
Giant Reed	<i>Phragmites australis</i>	X	X	
River Bulrush	<i>Bolboschoenus fluviatilis</i> (formerly <i>Scirpus fluviatillis</i>)			
Wild Celery ¹	<i>Vallisneria americana</i>		X	Yes
Shores and Waters to 0.25 m Deep		Identified in 1993	Identified in 2024	Recommended
Blue Flag	<i>Iris versicolor</i>			
Pickeralweed	<i>Pontederia cordata</i>			
Arrowhead	<i>Sagittaria latifolia</i>			Yes
Water Plantain	<i>Alisma plantago-aquatica</i>			
1. Not included in original 1993 recommendations. Planted in 2023.				

9.0 Recommendations

Burnside recommends the following for consideration to support aquatic life as Phase 2 rehabilitation begins:

Table 7: Summary of Recommendations to Improve Biological/Ecological Conditions During Phase 2

Biotic Group	Mechanism of Action	Effectiveness in Eutrophic Systems	Burnside Recommendations
Native Fish	Bio-manipulation via top-down control	Moderate – High minnow population may be suppressing desirable zooplankton.	Largemouth Bass stocking (>100 mm) may help reduce minnows if overwintering can be demonstrated.
Native Mussels	Nutrient sequestration	Low – many native mussels sensitive to poor water quality.	Paper Pondshell may provide some filtration, but unlikely to contribute significantly.
Zooplankton (esp. <i>Daphnia</i> spp.)	Grazing on phytoplankton	Moderate to high – if predation and chlorides are controlled.	Strong potential water quality improves and reduction in predation (i.e., increased habitat complexity and minnow control) Analysis of zooplankton community recommended before considering stocking
Macrophytes	Nutrient sequestration, habitat	High local effect; supports DO and stability.	Stabilizing effect once water quality permits growth
Habitat Complexity	Increase habitat complexity	High – Placement of woody material (stumps with root mass and elevated logs) or boulder complexes may enhance available habitat for Largemouth Bass.	Swan Lake currently offers limited structural habitat. Establishment of a bass population may be aided by increasing habitat complexity until macrophytes can re-establish. Deepening not recommended – evaluate after 2026 fish survey to determine bass survivability.

Nutrient and Oxygen Management

- Construction of bioswale may be most effective – low cost means to reduce and sustain nutrient levels and promote water circulation
- Enhance the existing bioswale to provide opportunities to beautify section of the lake
- Artificial aeration may benefit the lake (e.g., assistance in water circulation)

Chloride Reduction

- Maintain focus on source control (grit-separator rerouting, winter salt management, etc.)
- Target ≤ 120 mg/L Cl^- to aid restore zooplankton communities

Biological Recovery

- Zooplankton study may be useful to determine current species composition
- Add structural habitat (woody material, rock clusters) in absence of large macrophyte beds
- Determine if Largemouth Bass stocked in 2025 survive until 2026. If Largemouth bass can overwinter, they may help reduce minnow population to alleviate predation on zooplankton
- Delay further macrophyte plantings until turbidity is reduced

10.0 Conclusion

Success of Swan Lake's aquatic communities depends on improving water quality before attempting further biological restoration. Lack of noticeable improvements to the lake's water quality is likely a result of persistent problems interacting to keep lake conditions turbid.

Effective Recovery May Require

1. Nutrient stabilization through mechanical circulation via bioswale and/or artificial aeration
2. Chloride control by some means of reduction
3. Improvement to *Daphnia* spp. populations through improvements to habitat complexity, selective predator stocking, and eventual re-vegetation once clarity returns

This sequence offers a chance of achieving a self-sustaining mesotrophic system that supports a balanced, resilient, aquatic community.

Chloride-induced suppression of *Daphnia* spp. populations is likely exacerbating Swan Lake's phytoplankton blooms. Increasing *Daphnia* spp. abundance can yield short-term improvements (<2 years) in water clarity and phytoplankton control (Kibuye et al. 2021) and should be considered as part of broader lake-management planning. While some mechanism for chloride reduction is recommended, it alone will likely not resolve the issue. Creating a population of Largemouth Bass may help to improve the *Daphnia* spp. populations through the reduction of Fathead minnow. However, the current zooplankton community is unknown.

Improvement to the water quality of shallow, eutrophic lakes, such as the case in Swan Lake, are impeded by its slow recovery, as it is hard for people to maintain interest (May et al. 2020). However, actions for the improvements of Swan Lake are promising, and should encourage managers to maintain support and interest.

1.0 References

- Albright MF, Harman WN, Tibbits WT, Gray MS, Warner DM, Hamway RJ. 2004. Biomanipulation: a classic example in a shallow eutrophic pond. *Lake and Reservoir Management*. 20(4):263–269.
- Ali AM, Pickering NK. 2022. Systematic evaluation of materials to enhance soluble phosphorus removal using biofiltration or bioretention. *Water*. 14(21):3483. doi:10.3390/w14213483.
- Anderson P, Baldwin B. 2025. Swan Lake management options: literature review of management options and effects of increased oxygenation in Swan Lake [draft]. Project No. 3645. Waterloo (ON): Natural Resource Solutions Inc. 27 p.
- Bernes C, Carpenter SR, Gårdmark A, Larsson P, Persson L, Skov C, Speed JD, Van Donk E. 2015. What is the influence of a reduction of planktivorous and benthivorous fish on water quality in temperate eutrophic lakes? A systematic review. *Environmental Evidence*. 4(1):7. doi:10.1186/s13750-015-0032-9.
- Bunt CM, Peat TB, Cooke SJ. 2021. Seasonal movement and habitat use of Largemouth Bass in shallow Ontario lakes. *Environmental Biology of Fishes*. 104(2):165–179.
- Canadian Council of Ministers of the Environment (CCME). 2010. Canadian water quality guidelines for the protection of aquatic life: Ammonia (1999). In: Canadian environmental quality guidelines, 1999. Winnipeg (MB): CCME.
- Canadian Council of Ministers of the Environment (CCME). 2011. Canadian water quality guidelines for the protection of aquatic life: Chloride. In: Canadian environmental quality guidelines, 1999. Winnipeg (MB): CCME.
- Carpenter SR, Kitchell JF, Hodgson JR. 1987. Cascading trophic interactions and lake productivity. *BioScience*. 37(12):764–769.
- Czarnecka M. 2016. Coarse woody material as a shelter for fish in lowland rivers: importance of habitat complexity. *Hydrobiologia*. 767:21–33.
- Friends of Swan Lake Park (FOSLP). 2023. Appendix B: Recycling North Channel. In: Towards a comprehensive restoration plan for Swan Lake. Markham (ON): Friends of Swan Lake Park.
- Friends of Swan Lake Park (FOSLP). 2025a. Swan Lake Water Quality Plan: Summary of Phase 1 Discussions (2021–2025). Markham (ON): Friends of Swan Lake Park. 32 p.
- Friends of Swan Lake Park (FOSLP). 2025b. Swan Lake Water Quality Plan Review (Phase 1 Review Summary and Phase 2 Preparation). Markham (ON): Friends of Swan Lake Park. 33 p.

- Fullerton AH, Garvey JE, Wright RA, Stein RA. 2000. Overwinter growth and survival of largemouth bass: interactions among size, food, origin, and winter severity. *Transactions of the American Fisheries Society*. 129(1):1–12. doi:10.1577/1548-8659(2000)129<0001:OGASOL>2.0.CO;2.
- Geletu TT. 2023. Lake eutrophication: control of phytoplankton overgrowth and invasive aquatic weeds. *Lakes & Reservoirs: Research & Management*. 28(1):e12425.
- Godwin W, Coveney M, Lowe E, Battoe L. 2011. Improvements in water quality following biomanipulation of gizzard shad (*Dorosoma cepedianum*) in Lake Denham, Florida. *Lake and Reservoir Management*. 27(4):287–297. doi:10.1080/07438141.2011.609633.
- Ha JY, Saneyoshi M, Park HD, Toda H, Kitano S, Homma T, Shiina T, Moriyama Y, Chang KH, Hanazato T. 2013. Lake restoration by biomanipulation using piscivore and *Daphnia* stocking: results of the biomanipulation in Japan. *Limnology*. 14:19–30. doi:10.1007/s10201-012-0381-9.
- Hasler CT, Suski CD, Hanson KC, Cooke SJ, Tufts BL. 2009. The influence of dissolved oxygen on winter habitat selection by largemouth bass: an integration of field biotelemetry studies and laboratory experiments. *Physiological and Biochemical Zoology*. 82(2):143–152. doi:10.1086/591806.
- Hintz WD, Amott SE, Symons CC, Greco DA, McClymont A, Brentrup JA, et al. 2022. Current water quality guidelines across North America and Europe do not protect lakes from salinization. *Proc Natl Acad Sci U S A*. 119(9):e2115033119. doi:10.1073/pnas.2115033119.
- Keller AE, Ruessler DS. 1997. Determination or verification of host fish for nine species of unionid mussels. *Am Midl Nat*. 138(2):402–407.
- Kibuye R, Choi J, Choi MJ, An KG. 2021. Response of aquatic macrophytes and water quality to artificial aeration in small shallow lakes. *Water*. 13(2):134. doi:10.3390/w13020134.
- Lee J. 2019. Green infrastructure as a solution to hydrological problems: bioswales and created wetlands. *University of Florida Journal of Undergraduate Research*. 21(1).
- May L, Olszewska J, Gunn IDM, Meis S, Spears BM. 2020. Eutrophication and restoration in temperate lakes. *IOP Conference Series: Earth and Environmental Science*. 535:012001. doi:10.1088/1755-1315/535/1/012001.
- Metcalf-Smith JL, Staton SK, Mackie GL, Lane NM. 1998. Changes in the biodiversity of freshwater mussels in the Canadian waters of the lower Great Lakes drainage basin over the past 140 years. *J Great Lakes Res*. 24(4):845–858.

- Newman RM. 1991. Herbivory and detritivory in freshwater communities. *BioScience*. 41(9):633–641.
- Noda H, Maruyama A. 2014. The relationship between an introduced predator–prey pair, largemouth bass and bluegill, and the chlorophyll concentration in farm ponds. *Ichthyological Research*. 61(2):159–164. doi:10.1007/s10228-013-0380.
- Nohner JK, Taylor WW, Hayes DB, Roth BM. 2018. Influence of aquatic macrophytes on age-0 largemouth bass growth and diets. *Transactions of the American Fisheries Society*. 147(5):758–769. doi:10.1002/tafs.10067.
- OpenAI. 2025a. GPT-5 image enhancement [AI model]. San Francisco (CA): OpenAI. Available from: <https://openai.com/>.
- OpenAI. 2025b. GPT-5 image generation [AI model]. San Francisco (CA): OpenAI. Available from: <https://openai.com/>.
- Peat TB, Bunt CM, Cooke SJ. 2016. Winter habitat use by Largemouth Bass in small temperate lakes. *North American Journal of Fisheries Management*. 36(5):1012–1023.
- Piccioni M, Toffolon M, Lemaire BJ, Vinçon-Leite B. 2021. The thermal response of small and shallow lakes to climate change: new insights from 3D hindcast modelling. *Earth System Dynamics*. 12(2):439–455. doi:10.5194/esd-12-439-2021.
- Post DM, Kitchell JF, Hodgson JR. 1998. Interactions among adult demography, spawning date, growth rate, predation, overwinter mortality, and the recruitment of largemouth bass in a northern lake. *Canadian Journal of Fisheries and Aquatic Sciences*. 55(12):2588–2600. doi:10.1139/f98-139.
- Power ME. 1992. Top-down and bottom-up forces in food webs: do plants have primacy? *Ecology*. 73(3):733–746.
- Sarnelle O. 1993. Herbivore effects on phytoplankton succession in a eutrophic lake. *Ecological Monographs*. 63(2):129–149.
- Schaus MH, Godwin W, Battoe L, Coveney M, Lowe E, Roth R, Hawkins C, Vindigni M, Weinberg C, Zimmerman A. 2010. Impact of the removal of gizzard shad (*Dorosoma cepedianum*) on nutrient cycles in Lake Apopka, Florida. *Freshwater Biology*. 55(11):2401–2413. doi:10.1111/j.1365-2427.2010.02440.x.
- Scheffer M, Hosper SH, Meijer ML, Moss B, Jeppesen E. 1993. Alternative equilibria in shallow lakes. *Hydrobiologia*. 251(1):117–129.
- Scott WB, Crossman EJ. 1973. *Freshwater fishes of Canada*. Bulletin 184. Ottawa (ON): Fisheries Research Board of Canada.

- Setubal RB, Riccardi N. 2020. Long-term effects of fish biomanipulation and macrophyte management on zooplankton functional diversity and production in a temperate shallow lake. *Limnology*. 21(3):305–317.
- Siembida-Lösch B. 2021. Literature review of potential engineering solutions for the restoration of Swan Lake. Lindsay (ON): Fleming College Centre for Advancement of Water and Wastewater Technologies. Report prepared for Friends of Swan Lake Park. 23 p.
- Søndergaard M, Jensen JP, Jeppesen E. 2003. Role of sediment and internal loading of phosphorus in shallow lakes. *Hydrobiologia*. 506–509:135–145.
- Stahr KJ, Shoup DE. 2015. American water willow mediates survival and antipredator behavior of juvenile largemouth bass. *Transactions of the American Fisheries Society*. 144(5):903–910. doi:10.1080/00028487.2015.1052559.
- Toronto and Region Conservation Authority (TRCA). 2022. Shoreline restoration consulting project: community meeting 2 (Swan Lake) [PowerPoint presentation]. Toronto (ON): TRCA. 26 Apr 2022.
- Woodley A, Hintz LL, Wilmoth B, Hintz WD. 2023. Impacts of water hardness and road deicing salt on zooplankton survival and reproduction. *Scientific Reports*. 13:2975. doi:10.1038/s41598-023-30116-x.
- Zhang Y, Liu J, Chen Q, Wang H, Huang J, Zhou S. 2025. Eutrophication level constrains treatment efficacy of calcium peroxide in aquatic environments. *Environmental Advances*. 14:100520. doi:10.1016/j.envadv.2024.100520.
- Zhu S, Zhou W, Xu J, Chen M, Chen W. 2021. Role of benthic macroinvertebrates in the self-purification of lakes. *Water*. 13(10):1085. doi:10.3390/w13101085.

